

On The Electron Magnetic Moment Anomaly

André Michaud

SRP Inc Service de Recherche Pédagogique Québec Canada

Abstract:- It can be shown that the difference between the experimental value of the electron magnetic moment and that of the Bohr magneton is due to the magnetic drift of the electron's carrying energy induced at the hydrogen ground state gyroradius, corresponding to the electron being forced to move on a closed orbit about the nucleus in an isolated hydrogen atom.

Keywords:- Energy density, gyroradius, Bohr magneton, magnetic moment anomaly, circular motion, straight line motion, electron g factor, magnetic drift, first principles, atomic hydrogen atom.

I. STRAIGHT LINE MOTION AND EQUAL ELECTRIC AND MAGNETIC ENERGY DENSITY

Let us first summarize the various elements that must be considered to address the issue of the long standing still unexplained difference, seen as an anomaly, between the theoretically obtained Bohr magneton and the actual experimentally measured so-called "electron magnetic moment".

It was verified in a previous paper ([1], equation (35) and associated footnote) that equal local electric and magnetic energy densities imposes straight line motion to free moving localized photons and also of massive elementary particles, in conformity with Maxwell's theory.

The value of the Bohr magneton (μ_B) is obtained from the electron theoretical gyromagnetic moment equation, that is:

$$\frac{e}{m_0} = \frac{\mu_B}{S_z}, \quad \text{where } S_z = h/4\pi, \quad \text{so } \mu_B = \frac{eh}{4\pi m_0} = 9.27400899 \text{ E} - 24 \text{ J/T} \quad (1)$$

We also determined in paper ([1]) that the magnetic field of an energy quantum equal to that induced at the Bohr rest orbit ([1], equation (18)) involves exactly half of the energy induced at this distance from the nucleus ([1], equations (26) and (27), and associated footnote), which means that the so-called Bohr magneton (μ_B) **cannot possibly be a property of the electron proper**, but is rather a property of its carrying energy as induced at the Bohr gyroradius.

Now, the 3-spaces expanded Maxwellian space geometry ([3]) reveals that this magnetic energy can only oscillate at the rated energy frequency between an electric state and a magnetic state, a magnetic state whose moment can be experimentally measured.

For straight line motion, either as free photons or carrying-energy for massive particles, this cyclically oscillating energy will completely cross over from one state to the other at each cycle, which is what guarantees in this model equal local density of energy for both electric and magnetic aspects during each cycle, which in turn is the mandatory condition for straight line motion for free moving energy as shown from Maxwell's 4th equation, and also straight line motion for massive charged particles as shown by the Lorentz equation, respectively summarized by the two following equations:

$$c = \frac{E}{B} \text{ from Maxwell's 4}^{\text{th}} \text{ equation} \quad \text{and} \quad v = \frac{E}{B} \text{ from the Lorentz equation} \quad (2)$$

We also verified ([1], equation (33)) that the theoretical magnetic field of the energy quantum induced at the Bohr orbit can be calculated with equation:

$$B_0 = \frac{\mu_0 \pi e c}{\alpha^3 \lambda^2} \quad (3)$$

and that the relation between this magnetic field and the theoretical Bohr magneton is given by equation ([1], equation (25)):

$$B_0 = E / 2\mu_B = 235051.735 \text{ T} \quad (4)$$

where energy E is of course the Bohr rest orbit energy (4.35974377E-18 Joules) and the Bohr magneton is of course equal to 9.27400899E-24 J/T as calculated with equation (1).

II. CIRCULAR MOTION AND THE BOHR MAGNETON

The big question now is:

How is it that the Bohr magneton, theoretically meant to involve an electron **translating on a closed circular orbit** about the nucleus, can thus be equated to the magnetic field of a free electron **moving in straight line** with the same energy?

In 1909, Samuel Jackson Barnett ([4]) discovered that if a rod of demagnetized ferromagnetic material is suspended to a thin wire and made to rotate by any mechanical means, the rod becomes magnetized and that the intensity of the resulting macroscopic magnetic field proved to be directly proportional to the angular velocity of the rod as the velocity was caused to vary!

We analyze in a separate paper ([5]) that this could only be due to the unpaired electrons in the demagnetized rod to align orthogonally with respect to the rotation axis provided by the supporting thread, which causes the local magnetic field associated to the carrying energy that supports this circular translation motion of each of these electrons to align parallel and add up to become detectable by addition at the macroscopic level, and to logically intensify as their velocity, thus their energy, is increased.

III. CIRCULAR MOTION AND UNEQUAL FIELDS ENERGY DENSITY

On the other hand, it is well understood in circular accelerators circles ([6], p. 43) that when an electron is forced to move in a magnetic field that is not counterbalanced by an equal energy density electric field, it will start moving in a circle and if the magnetic field is increased further, the radius of that circle will diminish further.

The fundamental relativistic equation used in all existing closed circuit high energy accelerators, including the recently activated LHC, is the following:

$$qv\mathbf{B}_o = \gamma \frac{m_o v^2}{r_o} \quad (5)$$

From which the particle's magnetic orbit radius (named the gyroradius) equation is drawn:

$$r_o = \gamma \frac{m_o v}{q\mathbf{B}_o} \quad (6)$$

The Barnett effect effectively confirms that when electrons are forced to move in a circle, they will generate a magnetic field, which by definition will not be counterbalanced by an equal energy density electric field, since local equal energy density of electric and magnetic fields would mandatorily cause straight line motion of the charges involved, and that this magnetic field will increase as the electron's translation velocity (thus the associate carrying energy) is increased.

So why shouldn't the same Barnett effect also apply to a single electron forced into moving in a circle about an isolated proton (as in an isolated hydrogen atom)?

What do we know about the magnetic moment of the electron besides the Bohr magneton, which is calculated from theory (equation (1))? We know from experimental measurements that have been extensively carried out since the 1930's that the actual magnetic moment of the electron on the hydrogen rest orbital is measurably higher than the Bohr magneton!

What a surprise in light of all of these considerations, that the real magnetic moment of the electron in the hydrogen atom rest orbital would be higher than the Bohr magneton since we verified in paper ([1]) that the theoretically calculated Bohr magneton implicitly involves equal electric and magnetic energy densities associated to **straight line motion** of an electron having the same energy as the Bohr ground state energy, in blatant contradiction with the state of the actual electron captive in the rest state in an isolated hydrogen atom which is **forced to move in a circle!**

The difference between the Bohr magneton value and the experimental value is typically represented by a ratio of the latter over the former. The currently accepted value for this ratio, termed "**The electron magnetic moment anomaly**" ([2], p.1-3) is approximately:

$$\frac{\mu_e}{\mu_B} = 1.001159653 \quad (7)$$

which sets the current value ([2], p.1-3) of the experimentally verified electron magnetic moment to

$$\mu_e = 1.00115965 \ 3 \times \mu_B = 9.28476362 \ E - 24 \ J/T \quad (8)$$

IV. THE ELECTRON g FACTOR

The electron magnetic moment (μ_e) is currently calculated from the classical gyromagnetic moment equation previously mentioned (1) modified by the introduction of the **g factor of the electron**, whose

definition lies beyond the scope of this text, and whose value, theoretically set at 2, is further ad hoc finely corrected to $g/2 = 1.001159653$ to account as precisely as possible with the experimentally measured value of μ_e :

$$\mu_e = \frac{g}{2} \frac{eh}{4\pi m_o} = 9.28476362 \text{ E} - 24 \text{ J/T} \quad (9)$$

Note that this ratio is approximate to a certain extent since it can be measured only very indirectly and involves values for all hyperfine sub-states of the ground state of the hydrogen and deuterium atoms. For example, Julian Schwinger's 1947 paper on this issue ([7]) rates it at 1.001162.

More recently in paper ([8]), that is in 2006, the **g/2 factor** was established at 1.00115965218 with a different method. So any value in this range is likely to physically apply to one or other of the actual states or to a mean value of the ground level in the particular circumstances of measurement.

V. HIGHER LOCAL MAGNETIC ENERGY DENSITY FOR CIRCULAR MOTION

So this higher measured magnetic moment, coupled with the fact that the isolated hydrogen atom ground state electron can only move in a closed orbit, however uncertain its position may be at any given moment, obviously reveals that the magnetic field of the carrying energy of this electron will involve a magnetic carrying energy density higher than its electric carrying energy density since we know that equality would mandatorily involve straight line motion of the electron.

VI. LOWER CORRESPONDING LOCAL ELECTRIC FIELD FOR CIRCULAR MOTION

This means that the electron carrying energy magnetic field will be increased while its carrying energy electric field will be diminished in proportion to account for the physical fact that the electron is forced to move in a closed circle, while the mean total amount of carrying energy at the Bohr orbit has to remain invariant given that this total amount is dependent only on the mean distance to the nucleus.

VII. THE ELECTRON MAGNETIC DRIFT g FACTOR IS AN AD HOC QUANTITY

Let us emphasize here that the electron g factor, being an ad hoc quantity, **is not calculated from first principles**, since it is established only by comparing the experimentally measured electron magneton with the theoretical Bohr magneton. This in turn means that up to now, no theory has been able to link the observed magnetic drift associated with circular motion of elementary particles to first principles!

However, the 3-spaces model provides ample reasons to conclude that such forced stress drift of energy from electric state towards magnetic state of the carrying energy of massive particles captives of circular orbits is directly linked to, and varies with, the distance between the interacting charged particles. In other words, in accordance with observation, the tighter a closed circle an electron will be forced into, the more extensive will be the drift of its carrier-photon cycling energy from its electrical state towards its magnetic state.

VIII. CIRCULAR MOTION MAGNETIC DRIFT FROM FIRST PRINCIPLES

We will see presently that equation (33b) from previous paper ([9]) making use of distance related energy levels or equation (49) from the same paper making use of corresponding energy wavelengths allow calculating **from first principles** a value in the proper range without the need for any ad hoc corrective factor since both equations have been derived from first principles. Note that equation (55) derived from Special Relativity in paper ([1]) is identical to equation (49) derived from the 3-spaces model in paper ([9]).

Both equations (33b) and (49) from paper ([9]) in fact, allow calculating an effective energy drift ratio towards magnetic state for the whole range of possible interaction distances down to and including those of quarks up and down inside nucleons, thus possibly providing a direct theoretically derived method of explaining the drift from the unit charge of the electron to the partial charges of quarks up and down ([10], Chapter 17) and the drift ratio of their local carrying-photons, which could allow precise calculation of the observed magnetic moment of nucleons, as we are going to do in a coming paper.

Now why should these equations provide such a ratio?

Let us consider that they already provide a ratio of the actual relativistic velocity of a massive particle over the speed of light, calculated from the absolute wavelength of the energy related to the orbital distance of an electron about a hydrogen nucleus. Let us consider equation (49) from paper ([9]), for example, that we reproduce here:

$$\frac{v}{c} = \frac{\sqrt{\lambda_c(4\lambda + \lambda_c)}}{(2\lambda + \lambda_c)} \quad (10)$$

Where λ_c is the electron Compton wavelength and λ will be here the absolute wavelength of the electron carrying energy on the Bohr orbit.

Or equation (33b) from the same paper

$$\frac{v}{c} = \frac{\sqrt{4EK + K^2}}{2E + K} \quad (11)$$

Where E is the energy captive of the electron rest mass and K is the kinetic energy induced at the distance separating the electron orbit and the nucleus (here, the Bohr radius).

The direct relation of this velocity ratio to electron magnetic moment is that the velocity involved is the actual relativistic velocity that a free electron will have when moving in a straight line, when energy density is equal for both carrying electric and magnetic fields as explored in paper ([1]), when possessing an energy exactly equal to that induced at any given gyroradius about a hydrogen atom nucleus.

We will see now that dividing either of these equations by 2π to involve a relation normal to the direction of motion of the electron, as a gyroradius is with respect to the direction of motion of the electron on a circular orbit will provide a ratio in the exact range of the ad hoc electron g factor.

Let us now calculate the mean magnetic drift ratio for the hydrogen ground state with equation (10), where $\lambda_c = 2.426310215E-12$ m is the absolute wavelength of the rest mass energy of the electron and $\lambda = 4.556335256E-8$ m is the absolute wavelength of the energy induced in the electron at the Bohr radius (in this example, the Bohr gyroradius energy)

$$\text{magnetic_drift} = \frac{\delta\mu}{\mu_B} = \frac{\sqrt{\lambda_c(4\lambda + \lambda_c)}}{2\pi(2\lambda + \lambda_c)} = 1.161386535E-3 \quad (12)$$

By using the proper absolute carrying energy wavelength of each orbital that an electron can occupy in any atom, the proper magnetic drift ratio will be obtained for the particular gyroradius considered, which will allow calculating the pertaining local carrying energy's **drifted magnetic field** consistent with that orbital radius and the associated **reduced electric field** of this electron's carrying energy.

$$\mathbf{B}_d = \mathbf{B}_0 \times (1 + \text{magnetic_drift}) = 235324.3134 \text{ T} \quad (13)$$

In other words, as an electron's carrying energy oscillates between electric and magnetic states, part of it will be prevented by the stress due to closed circuit rotation from completely transferring to the electric state at each cycle. So that during each cycle, the mean energy making up the local magnetic field of the carrying energy will be equal to $(E/2) \times (1 + \text{magnetic drift})$ and the mean energy making up the corresponding local electric field will be $E - [(E/2) \times (1 + \text{magnetic drift})]$, this difference in the resulting mean energy density between local electric and magnetic states then directly explains the translation in circle of the electron at this particular gyroradius.

Alternately, the reciprocal equation ([11]) making use of energy can be used to cover the same complete range of possible magnetic drift ratios, where $E = 8.18710414 E-14$ Joules is the energy making up the rest mass of the electron and $K = 4.359743805 E-18$ Joules is the carrying energy of the electron (in this example, the Bohr gyroradius energy)

$$\text{magnetic_drift} = \frac{\delta\mu}{\mu_B} = \frac{\sqrt{4EK + K^2}}{2\pi(2E + K)} = 1.161386535E-3 \quad (14)$$

Now, the augmented drifted magnetic field (\mathbf{B}_d) from equation (13) that must physically exist at the Bohr radius due to the closed orbit circular motion involved is clearly seen to be increased beyond the value that it would have if the electron was moving in straight line with the same energy. Actually, this increased magnetic field is equal to that of a higher energy free photon or carrying photon that would be moving in straight line, in fact, a higher energy of which this magnetic field would constitute half the total complement when in straight line motion.

But since calculation of the corresponding electron magnetic moment (μ_e) requires using the energy corresponding to the increased drifted magnetic field and that this energy corresponds to half the energy of that higher energy photon, we need to calculate the energy of that higher energy photon before proceeding.

Equation (3) provides us with the key to this calculation since the only variable involved is the wavelength of the drifted magnetic field. So

$$\text{From } \mathbf{B}_d = \frac{\mu_0 \pi e c}{\alpha^3 \lambda^2} \text{ we define } \lambda = \sqrt{\frac{\mu_0 \pi e c}{\alpha^3 \mathbf{B}_d}} \quad (15)$$

And since $E = hc/\lambda$, we can write

$$E = h \sqrt{\frac{c \alpha^3 \mathbf{B}_d}{\mu_0 \pi e}} \quad (16)$$

So, from theory, and without using any ad hoc constant, we now obtain for the magnetic moment of the electron at the Bohr radius the following value:

$$\mu_{e_g} = \mu_B \times 1.00116138 \ 653 = 9.28477969 \ 4 \text{ E} - 24 \text{ J/T} \quad (17)$$

which stands barely outside the rated standard relative 4.0E-6 uncertainty factor of the measured value of 9.28476362 E-24 J/T.

As a final observation, we see that the phrase “**electron magnetic moment**” is quite a misnomer since its value specifically pertains to the specific mean energy of the electron hydrogen ground state carrying energy, that is, the energy of its carrier-photon, and should be named accordingly.

IX. CONCLUSIONS

There is thus ground to conclude that the so-called electron magnetic moment is only one discrete state of the range of all possible carrying energy magnetic moments that depend directly on the electron gyroradius within atoms.

REFERENCES

- [1]. André Michaud. **Field Equations for Localized Individual Photons and Relativistic Field Equations for Localized Moving Massive Particles**, International IFNA-ANS Journal, No. 2 (28), Vol. 13, 2007, p. 123-140, Kazan State University, Kazan, Russia. Also available at the following Internet location <http://www.gsjournal.net/Science-Journals/Essays/View/2257>.
- [2]. David R. Lide, Editor-in-chief. **CRC Handbook of Chemistry and Physics**. 84th Edition 2003-2004, CRC Press, New York. 2003.
- [3]. André Michaud. **The Expanded Maxwellian Space Geometry and the Photon Fundamental LC Equation**. International Journal of Engineering Research and Development e-ISSN: 2278-067X, p-ISSN: 2278-800X, www.ijerd.com Volume 6, Issue 8 (April 2013), PP. 31-45. (<http://ijerd.com/paper/vol6-issue8/G06083145.pdf>).
- [4]. S.J. Barnett. **Gyromagnetic and Electron-Inertia Effects**. Rev.Mod.Phys. Vol 7, 129 (1935).
- [5]. André Michaud. **On the Einstein-de Haas and Barnett Effects**. International Journal of Engineering Research and Development. e-ISSN: 2278-067X, p-ISSN: 2278-800X, www.ijerd.com Volume 6, Issue 12 (May 2013), PP. 07-11. (<http://ijerd.com/paper/vol6-issue12/B06120711.pdf>).
- [6]. Stanley Humphries, Jr. **Principles of Charged Particle Acceleration**, John Wiley & Sons, 1986.
- [7]. Julian Schwinger. On Quantum-electrodynamics and the Magnetic Moment of the Electron. Phys. Rev. 73, 416-417 (1948).
- [8]. B. Odom, D. Hanneke, B. D’Urso and G. Gabrielse. **New Measurement of the Electron Magnetic moment Using a One-Electron Quantum Cyclotron**. Phys. Rev. Let. 97, 030801 (2006).
- [9]. André Michaud. **From Classical to Relativistic Mechanics via Maxwell**. International Journal of Engineering Research and Development, e-ISSN: 2278-067X, p-ISSN: 2278-800X, www.ijerd.com. Volume 6, Issue 4 (March 2013), PP. 01-10 (<http://ijerd.com/paper/vol6-issue4/A06040110.swf>).
- [10]. André Michaud. **Expanded Maxwellian Geometry of Space**, 4th edition, SRP Books, 2004. (Available in many eBook formats at <https://www.smashwords.com/books/view/163704>).